

Breastfeeding the Preterm Baby



By Nancy Mohrbacher, IBCLC, Lactation Consultant, Ameda Products
Co-author of *Breastfeeding Made Simple* and *The Breastfeeding Answer Book*

WHY YOUR MILK MATTERS

When you give birth early, your milk is higher in many of the nutrients that your baby needs, such as protein and iron. Your milk contains live cells that help protect your baby against infection and disease and strengthen your baby's immune system. Your milk has enzymes to help your baby digest food, and its growth factors help your baby's gut mature. No other food can give your baby these things.

Research has found that preterm babies who miss out on mother's milk may get sick more often and more severely. Later on, they also may have more vision problems, bone problems, and delays in development.

BIRTH TO DAY 4

Your preterm baby may breastfeed well from birth. But if not, start pumping. The sooner, the better. If you can, start within the first six hours. Use a hospital-grade rental pump. Plan to double pump 8-10 times per day for 10-15 minutes. For more details, see our sheet, "How to Bring In a Full Milk Supply with a Breast Pump."

Expect to pump just a little milk at first. But even drops are important to your baby. Every pumping you do "puts in your order" for more milk. This is how you can tell your body to make more milk.

DAYS 4 TO 10

Expect to see a big change by Day 3 or 4. Where there were drops, you'll likely see ounces. If your baby is not yet breastfeeding, use your pump to try to bring in a full supply by Day 10. Your body is primed and ready to do this right after birth. If you wait until later, building a full supply can be much harder. When you reach 25 to 35 ounces (750 to 1050 ml) per day, you are there.

When a baby is born early, many mothers grieve the loss of the birth they had planned. Although your baby may need medical help, providing your milk is the one thing **only you** can do for your baby. Your milk can be key to your baby's health.



When you start to pump more milk on Day 3 or 4, try these tips to boost your supply faster:

- **Pump longer**—until 2 minutes after the last drop of milk or 20-30 minutes in total. (Drained breasts make milk faster.)
- **Don't go longer than 5 hours** between pumpings until you reach a full supply. (Full breasts make milk slower.)
- **Pump 8-10 times each 24 hours.** You don't have to pump on a set schedule. You can pump every hour for part of the day to fit them all in.
- **Massage your breasts** during pumping.
- **Check your breast flange fit.** Flange fit can change as you pump more. See the photos in "Getting a Good Flange Fit" under "Breast Pumping" on ameda.com

Once they have a full supply, most mothers can pump less—6-7 times per day—to maintain their supply. If you can sleep all night without too much breast fullness, go ahead. Any milk that your baby does not need can be frozen for later.

The following extra steps have been found to be of no benefit to you or your baby: cleaning the breasts before pumping, sterilizing pump parts

after each use, and throwing away the first milk pumped. Most hospitals no longer recommend these steps.

STORING YOUR MILK

Follow your hospital's rules for milk collection and storage. Most hospitals suggest that you:

- Use glass or plastic containers approved for food storage.
- Use a tight-fitting lid, rather than a bottle nipple.
- Label the milk with the baby's name and the date and time of the pumping.

Your hospital may give you bottles and lids to use.

Milk storage times may differ from those for full-term healthy babies. Ask your hospital for the storage times it recommends at room temperature, refrigerator, and freezer. You can use the Ameda Cool 'N Carry insulated tote to keep the milk cool between home and hospital.

BOOSTING MILK SUPPLY

If your supply drops to below 25 ounces (750 mL) per day, try these ideas to bring it up.

- Pump longer. (Drained breasts make milk faster.)
- Hold your baby skin-to-skin before pumping.
- Switch to a hospital-grade rental pump, if you're not already using one.
- Pump more times each day.
- Ask your lactation consultant to talk to your physician about prescribed medications and/or over-the-counter preparations that can boost milk supply.

STARTING TO BREASTFEED

Your baby's ability to breastfeed will depend on how early your baby was born and her health. Your hospital may also have policies that affect when you start breastfeeding. Even before then, try to hold your baby skin-to-skin as much as you can. This helps your baby stay warm and calm, and sleep better. It may also help you make more milk.

Your baby has feeding skills even before you start breastfeeding. Research done in Sweden, where 98% of mothers breastfeed, found that:

- At 28 weeks gestation, babies can root at the breast and latch-on,
- At 31 weeks gestation, some babies can suck and get milk.



YOUR MILK IS THE ONLY SOURCE OF CERTAIN LIVE CELLS AND ENZYMES THAT HELP PROTECT YOUR BABY'S HEALTH.

- By 36 weeks gestation, 85 percent of preterm babies can fully breastfeed.
- Babies with health problems tend to take longer to breastfeed well.

When ready to take food by mouth, babies can go to the breast first. Studies have found that preterm babies may have fewer heartbeat and breathing problems when fed by breast than by bottle. A baby who is not taking full feedings at the breast can be tube-fed until she is breastfeeding well.

When it is time to give the breast, start breastfeeding when your baby is calm and alert. To know if your baby is ready, look for signs like:

- Sucking around the gavage tube.
- Putting her hand to her mouth.
- Nuzzling and turning her head toward you with a wide open mouth.

Think of your first breastfeedings as practice.

- Your baby may lick or mouth the nipple at first.
- Many preterm babies suck in short bursts and fall asleep quickly.
- It may take several feedings before your baby breastfeeds well.
- If your baby does not get much milk at first, it's okay, because she will be given more milk after breastfeeding.

To make the move to full breastfeeding, seek help from a board-certified lactation consultant (IBCLC). To find one near you, check www.ilca.org.

This is general information and does not replace the advice of your healthcare provider. If you have a problem you cannot solve quickly, seek help right away.

Every baby is different, and your baby may not be average. If in doubt, contact your physician or other healthcare provider.